



TOBACCO USE AMONG AFRICAN AMERICANS

Each year, approximately 45,000 African Americans die from smoking-related disease.¹ Smoking-related illnesses are the number one cause of death in the African-American community, surpassing all other causes of death, including AIDS, homicide, diabetes, and accidents.²

Tobacco Use Among African-American Youth

According to the National Youth Tobacco Survey (NYTS), released by the U.S. Centers for Disease Control and Prevention (CDC) and the Food and Drug Administration (FDA), e-cigarettes are the most commonly used tobacco product among African-American youth. In 2023, 5.6% of African-American high schoolers and 5.7% of middle schoolers, were current e-cigarette users.³

Cigars are the second most popular product among African-American youth, after e-cigarettes. African-American youth smoke cigars at twice the rate of white youth.⁴

African-American high school students smoke cigarettes at a much lower rate than their Hispanic and White peers, due to protective factors such as greater price sensitivity, religiosity, negative community attitudes towards smoking, parental opposition and sports involvement.⁵ In 2020, 2.8 percent of African American high school students smoked cigarettes* compared to the 5.3 percent of Whites and 4.6 percent of Hispanics.⁶

However, disparities persist and reductions in adolescent combustible tobacco product use have not equitably reached communities of color, particularly Black youth. A study evaluating combustible tobacco product use among adolescents in the U.S. from 2014 to 2020 found that the odds of combustible tobacco smoking declined 21.5% per year among White youth, which was 4.22 times greater than for Black youth at 5.1% per year.⁷ Furthermore, as African American youth enter young adulthood, their smoking rates begin to mirror that of their White peers.⁸

Tobacco Use Among African-American Adults

In contrast to youth usage patterns, African American adult smoking rates are nearly equivalent to that of Whites. According to the 2021 National Health Interview Survey (NHIS) of adults ages 18 and over, 11.7 percent of African-American adults in the United States are current smokers, compared to 12.9 percent of Whites, 7.7 percent of Hispanics, and 5.4 percent of Asian Americans.[†] Overall, 11.5 percent of US adults are current smokers. Nearly one in five (18.1%) African American adults are current users of any tobacco product, and they have the highest prevalence of cigar smoking of all major American racial/ethnic groups. 5.1 percent of African American adults are current cigar smokers, compared to 3.5 percent of adults overall.⁹

Despite these high smoking rates, research has shown that African-American smokers are more likely to think that smoking is socially unacceptable and are highly motivated to quit.¹⁰ In fact, more than 70 percent of current African-American smokers want to quit, and more than 60 percent made a quit attempt in the previous year.¹¹

Even though African Americans tend to initiate smoking later in life and African American smokers are more likely than White smokers to have used cessation counseling services in the previous year, they are less likely than White smokers to successfully quit smoking.¹²

* NYTS 2023 did not report a rate for current cigarette smoking among Black high school students as it had in previous years.

† NHIS 2021 did not report a rate for current cigarette smoking among AI/AN adults as it had in previous years.

- Data from 2015 NHIS show that, among smokers who made a quit attempt in the past year, only 4.9 percent of African Americans remained abstinent after 6 months, compared to 7.1 percent of Whites.¹³
- Another survey found that while the cessation rate for early- and late-onset white smokers as well as early-onset African American smokers was between 52-57%, it was only 33% for late-onset African American smokers.¹⁴
- African Americans may have lower cessation rates than whites because African Americans generally have higher levels of nicotine dependence as a consequence of their preference for mentholated cigarettes.¹⁵
- Compared to Whites, African-Americans are less likely to receive advice to quit from a health professional and are less likely to use proven cessation medication treatments which may also contribute to lower cessation rates.¹⁶

Disproportionate Health Burden of Tobacco Among African Americans

According to the 2021 NHIS, 53.7% of African American adults who ever smoked cigarettes reported having quit.¹⁷ African Americans suffer tremendous tobacco-related morbidity and mortality despite the fact that they tend to smoke fewer cigarettes per day and begin smoking later in life.¹⁸ African Americans are less likely to quit smoking, which means that they ultimately smoke for a longer period of their life which, contributing to mortality disparities.¹⁹

Smoking is responsible for about thirty percent of all cancer deaths.²⁰ African Americans have the highest death rates and shortest survival of any other racial or ethnic group for most cancers.²¹ Each year, more than 72,000 African Americans are diagnosed with a tobacco-related cancer and more than 39,000 die from a tobacco-related cancer.²² Smoking is responsible for 81 percent of lung cancer deaths.²³

- Lung cancer is the second most common cancer in both African-American men and women, but it kills more African Americans than any other type of cancer.²⁴ In 2022, it was estimated that 25,690 African Americans will be diagnosed with lung cancer and 14,160 African Americans will die from it.²⁵

From 2007 to 2016, both the overall cancer and lung cancer-specific death rates for African Americans declined faster than for Whites, greatly reducing disparities, especially among men. These declines are the result of a larger decrease in smoking initiation among young African-Americans over the previous 40 years. In fact, the larger decline in smoking among African American youth have led to a convergence of lung cancer incidence overall and a near elimination of the racial disparity among those less than 50 years old. However, African-American men have a 10 percent higher incidence rate and 14 percent higher death rate from lung and bronchial cancers (2016-2020) compared to White men.²⁶

Smoking is also a major cause of heart disease and stroke — conditions that kill more people in the African-American community than lung cancer.²⁷ Compared to non-Hispanic Whites, African-Americans are 40% more likely to die from heart disease, 50% more likely to have a stroke, and 50% more likely to die from a stroke.²⁸ In addition, smoking is a major cause of chronic bronchitis, emphysema, gastric ulcers, and cancers of the mouth, pharynx, larynx, esophagus, pancreas, uterine cervix, kidney, stomach, liver, colon and bladder.²⁹ Smoking can also cause worse health outcomes in cancer patients and survivors.³⁰

African Americans are also disproportionately affected by exposure to secondhand smoke. More than a third (39.7%) of African Americans are still exposed to this preventable health hazard.³¹ Rates are even higher among youth. Among African-American kids aged three to 11 years, two-thirds (66.2%) are exposed to secondhand smoke, compared to 38.1 percent of white kids of the same age.³² Exposure to secondhand smoke is known to cause sudden infant death syndrome (SIDS), respiratory infections, ear infections and more severe asthma attacks in children, as well as heart disease, stroke and lung cancer in adults.³³

Impact of Disproportionate Use of Menthol Cigarettes Among African Americans

The higher rates of some tobacco-caused disease among African Americans than whites may result, in part, from the greater use of menthol cigarettes among African-American smokers.³⁴ Overall, 85% of

African-American smokers smoke menthol cigarettes, compared to 29% of white smokers.³⁵ The popularity of menthol is also evident in the cigarette brand preferences of African American youth who smoke. More than two-thirds (69.1%) of African-American youth ages 12-17 prefer Newport brand cigarettes. In contrast, the predominant brand among white youth who smoke is non-menthol Marlboro.³⁶ Data from a nationally representative cross-sectional study found that 44.5 percent of African American menthol smokers would quit smoking in response to a prohibition on menthol cigarettes.³⁷

In 2013, the U.S. Food and Drug Administration (FDA) released a report finding that menthol cigarettes lead to increased smoking initiation among youth and young adults, greater addiction, and decreased success in quitting smoking. The agency concluded that, “these findings, combined with the evidence indicating that menthol’s cooling and anesthetic properties can reduce the harshness of cigarette smoke and the evidence indicating that menthol cigarettes are marketed as a smoother alternative to nonmenthol cigarettes, make it likely that menthol cigarettes pose a public health risk above that seen with nonmenthol cigarettes.”³⁸ The Tobacco Products Scientific Advisory Committee (TPSAC), in its 2011 report to the FDA, estimated that by 2020, 4,700 excess deaths in the African American community will be attributable to menthol cigarettes, and over 460,000 African Americans will have started smoking because of menthol cigarettes.³⁹

Research published since FDA’s and TPSAC’s reports continue to bolster these findings:

- In 2020, a study analyzing four waves of data from the government’s Population Assessment of Tobacco and Health (PATH) study found that among daily smokers, menthol cigarette smokers have a 24% lower odds of quitting as compared to non-menthol smokers (the study did not find a significant difference among quit rates for non-daily menthol and non-menthol smokers). Among daily smokers, African American menthol smokers had a 53% lower odds of quitting compared to African American non-menthol smokers and white menthol smokers had a 22% lower odds of quitting compared to white non-menthol smokers.⁴⁰
- A meta-analysis of findings from nearly 150,000 smokers found that among African Americans, menthol smokers have a 12% lower odds of smoking cessation compared to non-menthol smokers.⁴¹
- A 2014 randomized clinical trial of FDA-approved cessation treatments among 1,500 US adult smokers found that menthol smoking was associated with reduced likelihood of quitting, compared to non-menthol smoking. African American female menthol smokers had the lowest quit rates of all groups in the study.⁴²
- Relying on these studies as well as the FDA’s and TPSAC’s findings, the 2020 Surgeon General Report on Smoking Cessation determined that the evidence was suggestive, but not conclusive as to the role of menthol on smoking cessation, finding the strongest evidence for reduced likelihood of smoking cessation among African American menthol smokers.⁴³ Due to the lower likelihood of smoking cessation among African American menthol smokers, the 2020 Surgeon General Report on Smoking Cessation concluded that, “Use of menthol cigarettes has been shown to contribute to tobacco cessation-related disparities in the United States.”⁴⁴

Reducing Tobacco Use among African Americans

Services and policies to help people quit using tobacco consist of a variety of evidence-based, individual and population-level approaches. According to the U.S. Public Health Service Clinical Practice Guideline, tobacco cessation treatments are effective across a broad range of populations. It is critical that health care providers screen for tobacco use and provide advice to quit to African American tobacco users.⁴⁵

Menthol cigarettes pose a tremendous public health threat, especially among Black Americans. In 2011, FDA’s Tobacco Products Scientific Advisory Committee (TPSAC) recommended that, “Removal of menthol cigarettes from the marketplace would benefit public health in the United States.”⁴⁶ Further, researchers estimate that a menthol ban would close the gap between lung cancer death rates for Black Americans and other U.S. racial and ethnic groups by 2025 – twenty-five years sooner than it otherwise

would have, and that by 2060, smoking attributable lung cancer death rates would be cut in half for Black Americans.⁴⁷

Implementation of public education campaigns reduce the number of youth who start smoking, increase the number of smokers who quit, and make tobacco industry marketing less effective, saving lives and health care dollars. Research from the 2013 *Tips From Former Smokers (Tips)* campaign shows that these campaigns can be particularly effective for and well-received by African Americans. The quit attempt rate for African American smokers in media markets with higher doses of the *Tips* campaign was 60 percent higher than that of smokers in standard dose media markets.⁴⁸

Policy interventions can also help people quit smoking. Research studies have established that cigarette price increases are particularly effective in preventing and reducing smoking among African Americans, youths, males, and persons in low-income households.⁴⁹ For example, a 10 percent increase in cigarette prices will reduce overall youth smoking by about seven percent but will reduce smoking among African-American male teenagers by 16 percent.⁵⁰ Although price increases are the most powerful anti-smoking factor among all youths, enforcing laws that prohibit sales of cigarettes to kids have also been found to be especially effective in reducing smoking among African-American teens.⁵¹

Finally, research indicates that African Americans have benefited less than other racial and ethnic groups from the spread of smokefree laws across the country, contributing to their disproportionate exposure to secondhand smoke.⁵² The 2012 Surgeon General's Report concluded that in combination with sustained programs and mass media campaigns, tax increases, and other regulatory initiatives, smokefree laws effectively reduce smoking among youth and young adults.⁵³

Campaign for Tobacco-Free Kids, January 26, 2024, Marela Minosa

More information on Tobacco and African Americans is available at
<https://www.tobaccofreekids.org/fact-sheets/tobaccos-toll-health-harms-and-cost/toll-of-tobacco-on-specific-populations-african-americans>

¹ US Department of Health and Human Services (HHS), "Tobacco Use Among US Racial/Ethnic Minority Groups—African Americans, American Indians and Alaskan Natives, Asian Americans and Pacific Islanders, and Hispanics: A Report of the Surgeon General," 1998, http://www.cdc.gov/tobacco/data_statistics/sgr/1998/complete_report/pdfs/complete_report.pdf.

² American Cancer Society, "Cancer Facts & Figures for African Americans, 2013–2014," 2013, <http://www.cancer.org/acs/groups/content/@epidemiologysurveillance/documents/document/acspc-036921.pdf>. American Heart Association, "African Americans and Cardiovascular Diseases: Statistical Fact Sheet, 2012 Update," http://www.heart.org/idc/groups/heart-public/@wcm/@sop/@smd/documents/downloadable/ucm_319568.pdf.

³ Birdsey J, et al. "Tobacco Product Use Among U.S. Middle and High School Students — National Youth Tobacco Survey, 2023." *MMWR* 72(44):1173–1182, November 3, 2023, <https://www.cdc.gov/mmwr/volumes/72/wr/pdfs/mm7244a1-H.pdf>.

⁴ Birdsey J, et al. "Tobacco Product Use Among U.S. Middle and High School Students — National Youth Tobacco Survey, 2023." *MMWR* 72(44):1173–1182, November 3, 2023, <https://www.cdc.gov/mmwr/volumes/72/wr/pdfs/mm7244a1-H.pdf>.

⁵ Garrett, BE, et al., "The African American Youth Smoking Experience: An Overview," *Nicotine & Tobacco Research*, 18(S1): S11-S15, 2016.

⁶ CDC, "Tobacco Product Use Among Middle and High School Students—United States, 2020," *MMWR*, 69, December 17, 2020, <https://www.cdc.gov/mmwr/volumes/69/wr/pdfs/mm6950a1-H.pdf>. Current smoker defined as having smoked in the past month.

⁷ Mantey, Dale S., et al. "Racial and Ethnic Disparities in Adolescent Combustible Tobacco Smoking from 2014 to 2020: Declines are lagging among non-Hispanic Black Youth." *Nicotine and Tobacco Research* (2024): ntae001.

⁸ Alexander, LA, et al., "Why we must continue to investigate menthol's role in the African American smoking paradox," *Nicotine & Tobacco Research*, 18(S1): S91-S101, 2016; ⁸ Roberts, ME, et al., "Understanding tobacco use onset among African Americans," *Nicotine & Tobacco Research*, 18(S1): S49-S56, 2016.

⁹ Cornelius ME, Loretan CG, Jamal A, et al. Tobacco Product Use Among Adults – United States, 2021. *MMWR Morb Mortal Wkly Rep* 2023;72:475–483. DOI: <http://dx.doi.org/10.15585/mmwr.mm7218a1>.

¹⁰ Royce, J, et al., "Smoking cessation factors among African Americans and Whites: COMMIT Research Group," *American Journal of Public Health* 83(2):220-6, February 1993.

¹¹ CDC, "Quitting Smoking Among Adults—United States, 2000-2015," *MMWR*, 65(52): 1457-1464, January 6, 2017, <https://www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6552a1.pdf>;

¹² CDC, "Current Cigarette Smoking Among Adults—United States, 2005-2015," *Morbidity & Mortality Weekly Report*, 65(44): 1205-1211, November 11, 2016, http://www.cdc.gov/mmwr/volumes/65/wr/mm6544a2.htm?s_cid=mm6544a2_w.

- ¹³ CDC, "Quitting Smoking Among Adults—United States, 2000-2015," *MMWR*, 65(52): 1457-1464, January 6, 2017, <https://www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6552a1.pdf>.
- ¹⁴ Roberts, ME, et al., "Understanding tobacco use onset among African Americans," *Nicotine & Tobacco Research*, 18(S1): S49-S56, 2016.
- ¹⁵ Royce, J, et al., "Smoking cessation factors among African Americans and Whites: COMMIT Research Group," *American Journal of Public Health* 83(2):220-6, February 1993.
- ¹⁶ CDC, "Quitting Smoking Among Adults—United States, 2000-2015," *MMWR*, 65(52): 1457-1464, January 6, 2017, <https://www.cdc.gov/mmwr/volumes/65/wr/pdfs/mm6552a1.pdf>; Giovino GA, et al. Differential Trends in Cigarette Smoking in the USA: Is Menthol Slowing Progress? *Tobacco Control*, August 30, 2013.
- ¹⁷ Cornelius ME, Loretan CG, Jamal A, et al. Tobacco Product Use Among Adults – United States, 2021. *MMWR Morb Mortal Wkly Rep* 2023;72:475–483. DOI: <http://dx.doi.org/10.15585/mmwr.mm7218a1>.
- ¹⁸ Alexander, LA, et al., "Why we must continue to investigate menthol's role in the African American smoking paradox," *Nicotine & Tobacco Research*, 18(S1): S91-S101, 2016; Roberts, ME, et al., "Understanding tobacco use onset among African Americans," *Nicotine & Tobacco Research*, 18(S1): S49-S56, 2016.
- ¹⁹ Holford, TR, et al., "Comparison of Smoking History Patterns Among African American and White Cohorts in the United States Born 1890 to 1990," *Nicotine & Tobacco Research*, 18(S1): S16-S29, 2016. See also, Kulak, JA, et al., "Differences in Quit Attempts and Cigarette Smoking Abstinences Between Whites and African Americans in the United States: Literature Review and Results from the International Tobacco Control US Survey," *Nicotine & Tobacco Research*, 18(S1):S79-S87, 2016.
- ²⁰ American Cancer Society. *Cancer Facts & Figures 2024*. Atlanta: American Cancer Society; 2024. <https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/2024-cancer-facts-figures.html>.
- ²¹ American Cancer Society. *Cancer Facts & Figures for African American/Black People 2022-2024*. Atlanta: American Cancer Society, 2022. <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/cancer-facts-and-figures-for-african-americans/2022-2024-cff-aa.pdf>
- ²² CDC, "Vital Signs: Disparities in Tobacco-Related Cancer Incidence and Mortality—United States, 2004-2013," *Morbidity & Mortality Weekly Report*, 65(44): 1212-1218, <http://www.cdc.gov/mmwr/volumes/65/wr/mm6544a3.htm>; See also American Cancer Society. *Cancer Facts & Figures for African American/Black People 2022-2024*. Atlanta: American Cancer Society, 2022.
- ²³ Siegel, R. L., et al. (2024). *Cancer statistics, 2024*. CA: a cancer journal for clinicians.; Islami F, et al. Proportion and number of cancer cases and deaths attributable to potentially modifiable risk factors in the United States. *CA: a cancer journal for clinicians*. 2018 Jan;68(1):31-54.
- ²⁴ American Cancer Society. *Cancer Facts & Figures 2024*. Atlanta: American Cancer Society; 2024. <https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/2024-cancer-facts-figures.html>.
- ²⁵ American Cancer Society. *Cancer Facts & Figures for African American/Black People 2022-2024*. Atlanta: American Cancer Society, 2022. <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/cancer-facts-and-figures-for-african-americans/2022-2024-cff-aa.pdf>.
- ²⁶ American Cancer Society. *Cancer Facts & Figures 2024*. Atlanta: American Cancer Society; 2024. <https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/2024-cancer-facts-figures.html>.
- ²⁷ American Cancer Society, "Cancer Facts & Figures for African Americans, 2019-2021," 2019, <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/cancer-facts-and-figures-for-african-americans/cancer-facts-and-figures-for-african-americans-2019-2021.pdf>.; Kochanek KD, et al. Deaths: Final data for 2020. *National Vital Statistics Reports*; vol 72 no 10. Hyattsville, MD: National Center for Health Statistics. 2023. DOI: <https://dx.doi.org/10.15620/cdc:131355>. See Table 8. *Number of deaths from selected causes, by Hispanic origin and race and sex: United States, 2020*.
- ²⁸ Kochanek KD, et al. Deaths: Final data for 2020. *National Vital Statistics Reports*; vol 72 no 10. Hyattsville, MD: National Center for Health Statistics. 2023. DOI: <https://dx.doi.org/10.15620/cdc:131355>. Table 10. *Age-adjusted death rate for selected causes, by Hispanic origin and race and sex: United States, 2020*.; Villarroel MA, et al. Tables of Summary Health Statistics for U.S. Adults: 2018 National Health Interview Survey. National Center for Health Statistics. 2019. Available from: <http://www.cdc.gov/nchs/nhis/SHS/tables.htm>.
- ²⁹ HHS, *The Health Consequences of Smoking: A Report of the Surgeon General*, 2004, http://www.cdc.gov/tobacco/data_statistics/sgr/2004/index.htm. HHS, *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General*, 2014, <http://www.surgeongeneral.gov/library/reports/50-years-of-progress/>.
- ³⁰ HHS, *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General*, 2014, <http://www.surgeongeneral.gov/library/reports/50-years-of-progress/>.
- ³¹ Brody DJ, Faust E, Tsai J. Secondhand smoke exposure among nonsmoking adults: United States, 2015–2018. *NCHS Data Brief*, no 396. Hyattsville, MD: National Center for Health Statistics. 2021. DOI: <https://doi.org/10.15620/cdc:101197>
- ³² Brody DJ, Lu Z, Tsai J. Secondhand smoke exposure among nonsmoking youth: United States, 2013–2016. *NCHS Data Brief*, no 348. Hyattsville, MD: National Center for Health Statistics. 2019. <https://www.cdc.gov/nchs/products/databriefs/db348.htm>
- ³³ HHS, *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General*, 2014, <http://www.surgeongeneral.gov/library/reports/50-years-of-progress/>.

- ³⁴ HHS, "Tobacco Use Among US Racial/Ethnic Minority Groups—African Americans, American Indians and Alaskan Natives, Asian Americans and Pacific Islanders, and Hispanics: A Report of the Surgeon General," 1998, http://www.cdc.gov/tobacco/data_statistics/sgr/1998/complete_report/pdfs/complete_report.pdf. HHS, *Reducing the Health Consequences of Smoking: 25 Years of Progress: A Report of the Surgeon General*, 1989, <http://sgreports.nlm.nih.gov/NN/B/B/X/S/>. See also Alexander, LA, et al., "Why we must continue to investigate menthol's role in the African American smoking paradox," *Nicotine & Tobacco Research*, 18(S1): S91-S101, 2016.
- ³⁵ Delnevo, CD, et al., "Banning Menthol Cigarettes: A Social Justice Issue Long Overdue," *Nicotine & Tobacco Research*, 22(10): 1673-1675, 2020.
- ³⁶ Substance Abuse and Mental Health Services Administration (SAMHSA)'s public online data analysis system (PDAS), National Survey on Drug Use and Health, 2021. https://datatools.samhsa.gov/#/survey/NSDUH-2021-DS0001/crosstab/?row=CIG30BR2&column=CATAG3&control=NEWTRACE2&weight=ANALWT_C&results_received=true.
- ³⁷ Pearson, JL, et al., "A Ban on Menthol Cigarettes: Impact on Public Opinion and Smokers' Intention to Quit," *American Journal of Public Health*, 102(11): e107-e114, 2012.
- ³⁸ FDA, "Preliminary Scientific Evaluation of the Possible Public Health Effects of Menthol Versus Nonmenthol Cigarettes," <http://www.fda.gov/downloads/ScienceResearch/SpecialTopics/PeerReviewofScientificInformationandAssessments/UCM361598.pdf>, 2013.
- ³⁹ Tobacco Products Scientific Advisory Committee, FDA, "Menthol Cigarettes and Public Health: Review of the Scientific Evidence and Recommendations, 2011, <http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/TobaccoProductsScientificAdvisoryCommittee/UCM269697.pdf>.
- ⁴⁰ Mills, S, et al., *The Relationship between Menthol Cigarette Use, Smoking Cessation and Relapse: Findings from Waves 1 to 4 of the Population Assessment of Tobacco and Health Study*, *Nicotine & Tobacco Research*, published online October 16, 2020, <https://doi.org/10.1093/ntr/ntaa212>.
- ⁴¹ Smith, PH, et al., "Use of Mentholated Cigarettes and Likelihood of Smoking Cessation in the United States: A Meta-Analysis," *Nicotine & Tobacco Research*, 2019, published online June 17, 2019.
- ⁴² Smith, SS, et al., "Smoking cessation in smokers who smoke menthol and non-menthol cigarettes," *Addiction*, 109: 2107-2117, 2014.
- ⁴³ U.S. Department of Health and Human Services. *Smoking Cessation. A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2020.
- ⁴⁴ U.S. Department of Health and Human Services. *Smoking Cessation. A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2020.
- ⁴⁵ Fiore, MC, et al., *Treating Tobacco Use and Dependence: 2008 Update—Clinical Practice Guideline*, U.S. Department of Health and Human Services. Public Health Service, May 2008, http://www.surgeongeneral.gov/tobacco/treating_tobacco_use08.pdf.
- ⁴⁶ Tobacco Products Scientific Advisory Committee (TPSAC), FDA, "Menthol Cigarettes and Public Health: Review of the Scientific Evidence and Recommendations, 2011, <https://wayback.archive-it.org/7993/20170405201731/https://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/TobaccoProductsScientificAdvisoryCommittee/UCM269697.pdf>
- ⁴⁷ Angelino, O, et al., "HowHow New Tobacco Control Laws Could Help Close the Racial Gap on U.S. Cancer," Council on Foreign Relations, February 1, 2023, <https://www.cfr.org/article/how-new-tobacco-control-laws-could-help-close-racial-gap-us-cancer>.
- ⁴⁸ McAfee, T, et al., "Increasing the dose of television advertising in a national antismoking media campaign: results from a randomised field trial," *Tobacco Control*, published online December 16, 2015.
- ⁴⁹ CDC, "Responses to Cigarette Prices By Race/Ethnicity, Income, and Age Groups – United States 1976-1993," *MMWR* 47(29):605-609, July 31, 1998. See also, Chaloupka, F & Pacula, F, "An Examination of Gender and Race Differences in Youth Smoking Responsiveness to Price and Tobacco Control Policies," *National Bureau of Economic Research*, Working Paper 6541, April 1998, www.uic.edu/~fjc/.
- ⁵⁰ Chaloupka, F & Pacula, F, "An Examination of Gender and Race Differences in Youth Smoking Responsiveness to Price and Tobacco Control Policies," *National Bureau of Economic Research*, Working Paper 6541, April 1998, www.uic.edu/~fjc/.
- ⁵¹ Chaloupka, F & Pacula, F, "An Examination of Gender and Race Differences in Youth Smoking Responsiveness to Price and Tobacco Control Policies," *National Bureau of Economic Research*, Working Paper 6541, April 1998, www.uic.edu/~fjc/.
- ⁵² Gonzalez, M, et al., "Strong Smoke-Free Law Coverage in the United States by Race/Ethnicity: 2000-2009," *American Journal of Public Health* 103(5): e62-e66, 2013.
- ⁵³ U.S. Department of Health and Human Services (HHS), *Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General*, HHS, U.S. Centers for Disease Control and Prevention (CDC), National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2012.